

B1b/S26

Fb1b/S26

FAO Fisheries Synopsis No. 26
(Distribution restricted)

F1b/S26 - Anchor
SAST-Anchovy - 1, 21(6), 002, 01

Provisional version

LIBRARY

JUN 4 1963

FISH & WILDLIFE SERVICE

SYNOPSIS OF BIOLOGICAL DATA ON ANCHOVY

Engraulis encrasicolus (Linnaeus) 1758

(Mediterranean and adjacent seas)

by

N. DEMIR

Hydrobiological Research Institute
Istanbul, Turkey

For

GENERAL FISHERIES COUNCIL
FOR THE MEDITERRANEAN

Library

NOAA, National Marine Fisheries Service
SWFC Honolulu Laboratory F/SWC2
2570 Dole Street
Honolulu, HI 96822-2396

FISHERIES DIVISION

FOOD AND AGRICULTURE ORGANIZATION OF THE UNITED NATIONS

Rome, February 1963

FAO
fish.
synop.
no. 26

3/B/30031p.

PURPOSE OF THIS SERIES

This is one of a series of synopses of biological data concerning species of aquatic organisms of economic importance. The primary purpose of this series is to make existing information readily available to fishery scientists, according to a standard pattern, and by so doing also to draw attention to gaps in knowledge. It is hoped that synopses in this series will be useful to scientists initiating investigations of the species concerned or of related ones; as a means of exchange of knowledge among those already working on the species, and as the basis for comparative study of fisheries resources.

The synopses are the output of a system for cumulative storage and retrieval of data on the biology of species and stocks, called Species and Stocks Thesaurus (SAST), maintained by the Fisheries Biology Branch of FAO. They will be brought up to date from time to time, as further information becomes available. It is, therefore, recommended that they be filed in loose-leaf folders; pagination begins anew with each chapter, so that partial revisions may be made.

Biology Branch is working to secure the cooperation of scientists in drafting synopses on species about which they have special knowledge, and welcomes offers of help in this task. Additions and corrections to synopses already issued will also be most welcome. Comments and requests for information should be addressed to the editor of this series:

Horacio Rosa, Jr.
Biology Branch, Fisheries Division
FAO, Rome, Italy

PREPARATION OF THIS SYNOPSIS

The first draft has been prepared for presentation at the 7th Session of the General Fisheries Council for the Mediterranean, Madrid, 12-18 March 1963.

It will be revised and republished as a synopsis of the same series.

Distribution

Author
7th Session of GFCM
FAO Fisheries Division
FAO Regional Fisheries Officers
Selectors 1 and 5

"Current Bibliography" entry

Demir, N. (1963) 8-01337 M
FAO Fish. Synops., (26):41p.
Synopsis of biological data on anchovy
Engraulis encrasicolus (Linnaeus) 1758
(Mediterranean and adjacent seas). Pro-
visional draft

Gives compiled information on identity, distribution, bionomics and life history, population, exploitation and a bibliography

C O N T E N T S

	<u>Page no.</u>
1 IDENTITY	1:1
1.1 <u>Taxonomy</u>	1
1.1.1 Definition	1
1.1.2 Description	1
- <u>Engraulis</u> Cuvier 1817	1
- <u>Engraulis encrasicolus</u> (Linnaeus 1758)	1
1.2 <u>Nomenclature</u>	2
1.2.1 Valid scientific name(s)	2
- <u>Engraulis encrasicolus</u> (Linnaeus) 1758	2
1.2.2 Synonyms	2
1.2.3 Standard common names, vernacular names	2
1.3 <u>General variability</u>	2
1.3.1 Subspecific fragmentation (races, varieties, hybrids)	2
2 DISTRIBUTION	2:1
2.1 <u>Delimitation of the total area of distribution and ecological characterization of this area</u>	1
2.2 <u>Differential distribution</u>	1
2.2.1 Areas occupied by eggs, larvae and other junior stages: annual variations in these patterns and seasonal variations for stages persisting over two or more seasons. Areas occupied by adult stages: seasonal and annual variations of these	1
2.3 <u>Behaviouristic and ecological determinants of the general limits of distribution and of the variations of these limits and of differential distribution</u>	1
3 BIONOMICS AND LIFE HISTORY	3:1
3.1 <u>Reproduction</u>	1
3.1.1 Sexuality (hermaphroditism, heterosexuality, intersexuality)	1
3.1.2 Maturity (age and size)	1
3.1.3 Mating (monogamous, polygamous, promiscuous)	1
3.1.4 Fertilization (internal, external)	1
3.1.5 Fecundity	1
- Relation of gonad size and egg number to body size and to age	1
- Coefficient of fecundity	1

	<u>Page no.</u>
3.1.6 Spawning	3:1
- Spawning seasons (beginning, end, peak)	1
- Number of spawnings per year, frequency	1
- Spawning time of day	6
- Induction of spawning, artificial fertilization	6
3.1.7 Spawning grounds	6
- Coastal (surface, vegetation, shore, shoal, sand, shelter); bottom	6
- Oceanic (surface, bottom)	6
3.1.8 Egg: structure, size, hatching type, parasites	6
- Structure and predators	6
- Size	6
- Parasites and predators	8
3.2 <u>Larval history</u>	8
3.2.1 Account of embryonic and juvenile life (prelarva, larva, postlarva, juvenile)	8
- Prelarva	8
- Larva, postlarva	8
- Juvenile	8
- Feeding	8
- Rates of: development and survival	8
- Periods of: development and survival	11
- Parental care	11
- Parasites and predators	11
3.3 <u>Adult history</u>	12
3.3.1 Longevity	12
3.3.2 Hardiness	12
3.3.3 Competitors	12
3.3.4 Predators	12
3.3.5 Parasites and diseases	12
3.3.6 Greatest size	12
3.4 <u>Nutrition and growth</u>	13
3.4.1 Feeding (time, place, manner, season)	13
3.4.2 Food (type, volume)	13
3.4.3 Relative and absolute growth patterns and rates	13
3.4.4 Relation of growth to feeding, to other activities, and to environmental factors*	13
3.5 <u>Behaviur</u>	13
3.5.1 Migration and local movements	13
3.5.2 Schooling	15
3.5.3 Reproductive habits	15

Page no.

4	POPULATION (STOCK)	
4.1	<u>Structure</u>	4:1
4.1.1	Sex ratio	1
4.1.2	Age composition	1
4.1.3	Size composition	1
4.2	<u>Size and density*</u>	
4.3	<u>Natality and recruitment*</u>	
4.4	<u>Mortality, morbidity</u>	2
4.4.1	Rates of mortality	2
4.4.2	Factors or conditions affecting mortality*	
4.4.3	Factors or conditions affecting morbidity*	
4.4.4	Relation of morbidity to mortality rates*	
4.5	<u>Dynamics of population*</u>	
4.6	<u>Relation of population to community and ecosystem, biological production, etc.</u>	2
5	EXPLOITATION	5:1
5.1	<u>Fishing equipment</u>	1
5.1.1	Fishing gear	1
5.1.2	Fishing boats	1
5.2	<u>Fishing areas</u>	1
5.2.1	General geographic distribution	1
5.2.2	Geographical ranges (latitudes, distances from coast, etc.)	1
5.2.3	Depth ranges	1
5.3	<u>Fishing seasons</u>	1
5.3.1	General pattern of fishing season*	
5.3.2	Duration of fishing season*	
5.3.3	Dates of beginning, peak and end of season	1
5.3.4	Variation in time or duration of fishing season*	
5.3.5	Factors affecting fishing season	1
5.4	<u>Fishing operations and results</u>	1
5.4.1	Effort and intensity	1
5.4.2	Selectivity*	
5.4.3	Catches	2
5.4.4	Past and present factors of effect on operations and results*	
5.5	<u>Fisheries management and regulations*</u>	
5.6	<u>Fish farming, transplanting and other intervention*</u>	

* As no information was available to the author, these items are not listed in the text.

1. IDENTITY

1.1 Taxonomy

1.1.1 Definition

Phylum VERTEBRATA

Subphylum Craniata

Superclass Gnathostomata

Series Pisces

Class Teleostomi

Subclass Actinopterygii

Order Clupeiformes

Suborder Clupeoidei

Superfamily Clupeoidae

Family Engraulidae

Genus Engraulis Cuvier 1817Species Engraulis encrasicolus (Linnaeus), 1758

1.1.2 Description

- Engraulis Cuvier 1817

Body elongate, barely compressed; cycloid scales moderate or large, thin and deciduous; no armed abdominal keel; snout conic, project beyond the lower jaw; eye large, without adipose eyelid; mouth extremely large; maxillary long, slender attached to cheek by a scarcely distensible membrane, extends back to mandible base or beyond; small teeth in jaws, on vomer, palatines and pterygoids; union of gill membranes short, leaves isthmus uncovered; branchiostegals short, vary from 9 to 14 in number; pseudobranchiae present; gill rakers long, slender, usually numerous; dorsal inserted about midway in body before anal and behind ventral origin.

- Engraulis encrasicolus (Linnaeus) 1758 (see Fig. 1)

Body rather slender; lower profile slightly more convex; head quite large, compressed; snout as long as broad; interorbital broadly convex; maxillary slender, extending backward far behind eye but not quite reaching preopercle edge; row of small teeth in jaws, along maxillary and mandible their whole extent; tongue finely granular; nostrils close to each other, hind one larger; dorsal short, high; anal longer and lower than dorsal; pectoral small; four oblong scales on upper caudal lobe, three on lower.

Color in life from greenish-blue to dark blue above, silvery white below.

Depth 7 to $7\frac{3}{5}$ in standard length; head 4 to $4\frac{4}{5}$; preopercle distance $2\frac{1}{10}$ to $2\frac{4}{10}$; predorsal distance $1\frac{9}{10}$ to $2\frac{1}{10}$; preanal distance $1\frac{4}{10}$ to $1\frac{5}{10}$; eye diameter 3 to $3\frac{7}{10}$ in head.

D. 14 to 17; A. 16 to 20; C. $n+19+n$; P. 14 to 19; V. 7; gill rakers 58 to 78- (26 to 35)+(32 to 43); Vertebrae 42 to 47-(24 to 27)+(18 to 20).

Described above from the Black, Marmara, Aegean Seas, Turkey examples, 79 to 144 mm total length.

The meristic counts given by various authors are shown in Table I.

1.2 Nomenclature

1.2.1 Valid scientific name(s)

Engraulis encrasicolus (Linnaeus) 1758

1.2.2 Synonyms

Clupea encrasicolus Linnaeus 1758
Clupea encrasicholus Lacépède 1798-1803
Clupea engraticolus Asso 1801
Clupea encrassicholus Delaroche 1809
Clupea engrasicholus Ramis 1814
Engraulis encrasicholus Cuvier 1817
Engraulis meletta Cuvier
Encrasicholus encrasicholus Fleming 1828
Engraulis vulgaris Nilsson 1832
Engraulis encrassicholus Greals 1855
Clupea encraciculus Poggi 1881

1.2.3 Standard common names, vernacular names.

See Table II.

1.3 General variability

1.3.1 Subspecific fragmentation (races, varieties, hybrids)

- Races

Although some statements have been made on the race composition of the Mediterranean anchovy, it is still an open question, as neither proportional measurements of body nor the mean numbers of the meristic counts (see Tables IV and V) give clear evidence for the separation of geographical races or subspecies. Furthermore, the influences of the environmental factors on the mean numbers of the meristic counts have not been taken into consideration by those authors who have studied this subject up to now. Fage (1911, 1920), however, maintains that the Mediterranean anchovy is a different race from the Atlantic anchovy, differing both morphologically and biologically. Hoek (1912), quoted by De Buen (1931) states that the Naples and Zuider Zee anchovies differ in body proportions.

Fage (1920) records the existence of two groups or local races of the Mediterranean anchovy: the eastern group in the Adriatic and Black Seas with 45.13 ± 0.155 mean number of vertebrae and the western group, spreading up to Gibraltar with 45.76 ± 0.275 as the mean number of vertebrae. Alexandrov (1927) suggests that the Mediterranean, Black and Azov Seas anchovies are three different races: 1. Engraulis encrasicolus mediterraneus Fage, in the Western Mediterranean and Adriatic, 2. Engraulis encrasicolus ponticus Alexandrov in the Western Black Sea, 3. Engraulis encrasicolus moeoticus Pusanov in the Eastern parts of the Azov and Black Seas. The characteristics of these races are shown in Table III.

Table I

Meristic counts of Engraulis encrasicolus

Author	D.	A.	C.	P.	V.	Br.	Gill rak.	Vert.
Günther, 1868	16-17	16-17	----	---	-	12-13	---	46-47
Moreau, 1881	15-18	16-18	2/21-22/3	15-17	7	12-13	---	---
Carus, 1889-1893	15-18	16-18	2/21-22/3	---	-	12-13	---	46-47
Kyle and Ehrenbaum, 1927	16-18	17-18	x+19+x	16-17	7	---	---	46-47
Fage, 1930	15-17	17-18	2/21	15-17	7	12-13	---	44-49
Fowler, 1936			----	---	-	---	30+30	---
Lübbert and Ehrenbaum, 1936	17-18	17-18	----	17	7	---	---	44-48
Lozano Rey, 1947	3+13-15	2+13-16	----	14-17	7	---	---	---
Berg et al., 1949	12-15	16-20	----	---	7	---	58-77	43-47
Căraușu, 1952	12-15	16-20	----	---	-	---	58-77	43-47(48)
Dieuzeide et al., 1954	15-18	16-18	2/21-22/3	15-17	7	---	---	44-49
Slastenenko, 1956	12-15	17	----	---	-	---	60-77	39-47
Popiel, 1962	16-18	17-18	---	16-17	7	---	---	46-47

Table II

Common and Vernacular Names

Country	Standard Common Name	Vernacular Name(s)
Algeria	Anchouba	Ancioja, Anchois, Anchoa, Boqueron
Bulgaria	Hamsia	Sardelka
Egypt	Anchooga	
France	Anchois	Antchoyo, Anchoye, Anchoia, Anchiove, Anchiuva, anxova, Amplova, Amplovo, Amploveto, Amplovino, Amplona, Amploua, Amarou, Ladrot
Greece	Gávros	Kapsia, Kapsi, Khampi, Chafia, Aphyi
Israel	Afian	
Italy	Acciuga	Alice, Alici, Aléce, Aoiccia, Alice'e sperone, Aliciastra, Aliciastruni, Ancioa, Anciona, Ancioja, Anciora, Anciova, Anchio, Anciva, Angioja, Anciovillo, Angiouvitta, Aggiuva, Azzuva, Amarou, Amplona, Amplova, Amplouva de fount, Bianco mangiare, Masculina, Masculinella, Masculino, Mascolini, Mezz'alici, Nini, Sardela, Sardone, Sardòn
Malta	Incova	Inciva, Inciova, Anciova, Nemusa
Monaco	Ancium	Ancina
Rumania	Hamsia	Amsi
Spain	Anchoa	Boquerón, Boquero, Bocartá, Bocarte, Bucaro, Anchova, Anxova, Aladroch, Sardinéta, Seito, Eledroc, Golfás
Tunisia	Anchouba	Anchouva
Turkey	Hamsi	
U.S.S.R.	Hamsa	Anchoye, Antschous
Yugoslavia	Brgljun	Brgljica, Kurcilj, Minčion, Inčun, Minčun

Table III

Characteristics of the races of Engraulis encrasicolus according to Alexandrov (1927)

Race	Average length (LF) in mm	$\frac{LL \times 100}{LF}$	$\frac{LD \times 100}{LF}$	$\frac{LA \times 100}{LF}$	D. rays	Vert
<u>Mediterranean</u>	125.0	21.72	44.00	58.15	12 - 14 13.20	44 - 48 45.44
<u>Ponticus</u>	118.0	21.37	41.76	57.64	12 - 15 13.70	43 - 47 44.62
<u>Maoticus</u>	98.9	23.08	44.72	60.35	12 - 15 13.40	39 - 46 43.57

· Majarova (1939), quoted by Slastenenko (1956) and Majarova-Chugunova (1954) distinguish two forms or natioes of the Black Sea anchovy: the eastern form Engraulis encrasicolus ponticus orientalis and the western form Engraulis encrasicolus ponticus occidentalis, each having a different mean number of vertebrae, dorsal rays, gill rakers, a different proportion of body, even different winter quarters, migration routes, spawning and feeding areas (see Tables IV, V and Section 3.5.1).

· Dulzetto (1938) considers that the anchovy of the Gulf of Catania is different from the stocks of the Black and Adriatic seas, Balearic Islands and Gulf of Lion even from the anchovies of the other Italian coasts. Vidalis (1949) believes that the anchovy of the Greek waters all belong to the same race.

The frequency distribution of the meristic counts given by various authors are shown in Table IV and V.

Table IV

Vertebral frequency distribution of Engraulis encrasicolus

Locality		Author	Number of fish having vertebral counts of:										N	M	S.D.	S.E.	Fl.M.
			39	40	41	42	43	44	45	46	47	48					
AZOV SEA		Alexandrov (1927)	1	2	4	20	86	118	33	2	-	-	266	43.57	-	-	0.063
BLACK SEA	Sevastopol	Fage (1911)	-	-	-	-	-	5	18	2	-	-	25	44.88	0.279	0.037	0.185
	St. 172	Fage (1920)	-	-	-	-	-	6	27	10	-	-	43	45.09	0.609	0.063	0.315
	Soviet waters	Alexandrov (1927)	-	-	-	-	4	55	49	8	1	-	117	44.55	-	-	-
	Eastern part	Majarova (1933) quoted by Slastenenko (1956)?	-	-	-	-	-	-	-	-	-	-	-	44.49	-	-	-
	Western part	Majarova (1939) quoted by Slastenenko (1956)?	-	-	-	-	-	-	-	-	-	-	-	44.45	-	-	-
	Bulgarian coast in 1952	Stoianov (1961)	-	-	-	2	15	139	422	193	5	-	776	45.03	-	-	-
	Bulgarian coast in 1954-55	Stoianov (1961)	-	-	-	-	21	315	424	32	-	-	792	44.59	-	-	-
	Fatsa Bay	Åasen and Akyüz(1956)	-	-	-	-	7	30	39	5	-	-	81	44.52	-	-	-
	Turkish coast in 1961	Demir (unpublished)	-	-	-	2	25	422	793	172	6	-	1420	44.82	0.6976	0.0185	0.0624
	SEA OF MARMARA (Gulf of Gemlik-Erdek) in 1961-62		Demir (unpublished)	-	-	-	-	6	174	795	168	9	-	1152	45.00	0.5909	0.017
AEGEAN SEA	Gulf of Izmir in 1961	Demir (unpublished)	-	-	-	-	2	69	320	136	5	-	532	45.13	0.6481	0.0229	0.0776
	Gulf of Thermai in 1948	Vidalis (1949)	-	-	-	-	-	-	-	-	-	-	45	44.53	0.9800	0.0940	0.4700
	Gulf of Euboea in 1948	Vidalis (1949)	-	-	-	-	-	-	-	-	-	-	360	45.31	0.6700	0.0230	0.1150
	Saronic Gulf in 1948	Vidalis (1949)	-	-	-	-	-	-	-	-	-	-	1143	44.90	0.5605	0.0110	0.0550
IONIAN SEA	Gulf of Patras in 1948	Vidalis (1949)	-	-	-	-	-	-	-	-	-	-	24	45.83	0.5900	0.0810	0.4050
	Gulf of Catania in 1936-37	Dulzetto (1938)	-	-	-	-	-	2	106	143	49	-	300	45.79	0.7108	0.0276	0.1380
ADRIATIC	Commachio	Fage (1920)	-	-	-	-	-	9	69	29	1	-	108	45.20	0.583	0.038	0.190
	Trieste	Giudice (1922) quoted by Dulzetto (1938)	-	-	-	-	-	-	-	-	-	-	-	45.37	-	-	-
	Venice	Giudice (1922) quoted by Dulzetto (1938)	-	-	-	-	-	-	-	-	-	-	-	44.11	-	-	-
TYRRHENIAN SEA	Cannitello	Giudice (1922) quoted by Dulzetto (1938)	-	-	-	-	-	-	-	-	-	-	-	45.02	-	-	-
	Spadofora	Giudice (1922) quoted by Dulzetto (1938)	-	-	-	-	-	-	-	-	-	-	-	45.20	-	-	-
	Gulf of Naples	Giudice (1922) quoted by Dulzetto (1938)	-	-	-	-	-	-	-	-	-	-	-	45.29	-	-	-
	Ischia	Giudice (1922) quoted by Dulzetto (1938)	-	-	-	-	-	-	-	-	-	-	-	44.36	-	-	-
LIGURIAN SEA	Portofino	Giudice (1922) quoted by Dulzetto (1938)	-	-	-	-	-	-	-	-	-	-	-	45.56	-	-	-
	Cornigliano	Giudice (1922) quoted by Dulzetto (1938)	-	-	-	-	-	-	-	-	-	-	-	45.31	-	-	-
Gulf of Lion		Fage (1911)	-	-	-	-	-	1	19	28	1	1	50	45.64	0.746	0.071	0.355
SPANISH COAST	Castellón coast in 1951-55	Larraneta (1958)	-	-	-	-	-	-	-	-	-	-	1249	45.67	0.630	-	0.05
	Almeria	Fage (1920)	-	-	-	-	-	-	7	17	6	-	30	45.96	0.655	0.081	0.405
	Málaga	Crehuat and Val-Cordón (1960)	-	-	-	-	-	-	112	464	91	-	667	45.97	-	-	0.0715

Table V

Dorsal fin-ray frequency distribution of *Engraulis encrasicolus*

Locality		Author	Number of fish having dorsal fin-ray counts of:						N	M	S.D.	S.E.	Fl.M.
			12	13	14	15	16	17					
AZOV SEA		Alexandrov (1927)	17	24	40	6	-	-	87	13.40	0.88	0.09	-
BLACK SEA	Sevastopol	Fage (1911)	-	7	15	1	-	-	23	13.1	-	-	-
	Soviet waters	Alexandrov (1927)	2	47	78	10	-	-	137	13.70	0.63	0.05	-
	Eastern part	Majarova (1939) quoted by Slastenenko (1956)	-	-	-	-	-	-	-	13.37	-	-	-
	Western part	Majarova (1939) quoted by Slastenenko (1956)?	-	-	-	-	-	-	-	13.08	-	-	-
	Turkish coast in 1961	Demir (unpublished)	-	-	2	59	74	5	140	15.59	0.588	0.0497	0.17
SEA OF MARMARA (Gulf of Gemlik-Erdek in 1961)		Demir (unpublished)	-	-	2	59	127	14	202	15.76	0.586	0.0413	0.14
AEGEAN SEA	Gulf of Izmir in 1961	Demir (unpublished)	-	-	2	53	46	1	102	15.45	0.559	0.0554	0.19
	Gulf of Thermai in 1948	Vidalis (1949)	-	-	-	-	-	-	94	15.17	0.730	0.0520	0.26
	Gulf of Euboea in 1948	Vidalis (1949)	-	-	-	-	-	-	373	14.67	0.680	0.0230	0.115
	Saronic Gulf in 1948	Vidalis (1949)	-	-	-	-	-	-	1799	14.52	0.650	0.0100	0.05
IONIAN SEA (Gulf of Patras) in 1948		Vidalis (1949)	-	-	-	-	-	-	26	15.32	0.540	0.073	0.367
Gulf of Catania in 1936-37		Dulzetto (1938)	-	2	165	133	-	-	300	14.43	0.509	0.0198	0.099
ADRIATIC	Trieste	Giudice (1922) quoted by Dulzetto (1938)	-	-	-	-	-	-	-	15.24	-	-	-
	Venice	Giudice (1922) quoted by Dulzetto (1938)	-	-	-	-	-	-	-	15.23	-	-	-
TYRRHENIAN SEA	Cannitello	Giudice (1922) quoted by Dulzetto (1938)	-	-	-	-	-	-	-	15.27	-	-	-
	Spadofoza	Giudice (1922) quoted by Dulzetto (1938)	-	-	-	-	-	-	-	15.35	-	-	-
	Gulf of Naples	Giudice (1922) quoted by Dulzetto (1938)	-	-	-	-	-	-	-	15.19	-	-	-
	Ischia	Giudice (1922) quoted by Dulzetto (1938)	-	-	-	-	-	-	-	14.11	-	-	-
LIGURIAN SEA	Portofino	Giudice (1922) quoted by Dulzetto (1938)	-	-	-	-	-	-	-	14.34	-	-	-
	Cornigliano	Giudice (1922) quoted by Dulzetto (1938)	-	-	-	-	-	-	-	14.32	-	-	-
Gulf of Lion		Fage (1911)	2	30	18	-	-	-	50	13.3	-	-	-
Balearic Islands		Fage (1911)	3	19	2	-	-	-	24	12.9	-	-	-

2. DISTRIBUTION

2.1 Delimitation of the total area of distribution and ecological characterization of this area

The Mediterranean and adjacent seas.

Ecological characters of this area vary greatly, specially from the point of view of temperature (annual average isotherms = 10 to 22°C) and salinity (5 to 40.0 ‰).

2.2 Differential distribution

2.2.1 Areas occupied by eggs, larvae and other junior stages: annual variations in these patterns and seasonal variations for stages persisting over two or more seasons. Areas occupied by adult stages: seasonal and annual variations of these

- Eggs

Eggs are distributed all over the Sea of Azov with the exception of the areas having lowest salinity (Nikolski, 1957); all over the Black Sea (Dekhnik, 1954; Vodjanitskii-Kazanova, 1954; Einersson-Gürtürk, 1960; Guéorghiev et al., 1961); Sea of Marmara except the parts deeper than 300 m. (Arim, 1957) and throughout all the other adjacent seas and the Mediterranean except for the central parts of the basins. Eggs become scarce along the coasts of Libya, Egypt, Israel and Southern Anatolia (Fage, 1920; De Buen, 1931).

- Larval stages

Larvae are found in all egg areas.

- Junior stages

Junior stages occur in all adult areas. Berg et al. (1949), however state that the juniors, unlike the adults, can stay in the shallow waters during the year. According to Fage (1920) these are only the juniors which are hatched in Autumn.

- Adult stages

See 2.1 and 3.5.1

2.3 Behaviouristic and ecological determinants of the general limits of distribution and of the variations of these limits and of differential distribution.

Since anchovy withstand 5 to 41.55 ‰ salinity and 6 to 29°C temperature ranges, there is no limiting factor for the general distribution in the Mediterranean and adjacent seas. But the eggs and larvae are not found in waters which have a temperature lower than 13°C (the most favorable temperature range being 20 to 25°C; salinity 9 to 37.5‰).

In Spring, the occurrence of the adults in the surface layers of coastal waters depend mainly on the surface temperature of the water and the abundance of plankton as food. In Autumn, the migration to the deeper layers also depends mainly on the temperature. When the surface temperature of the waters drops below 6 or 7°C, the anchovy pass into deeper and warmer layers.

3. BIONOMICS AND LIFE HISTORY

3.1 Reproduction

3.1.1 Sexuality (hermaphroditism, heterosexuality, intersexuality)

Heterosexual without external secondary characters. The case of hermaphroditism has been observed by Mužinić (1956) in one specimen from the Adriatic.

3.1.2 Maturity (age and size)

Anchovy reach maturity at the end of their first year of life. Fage (1920) states that the individuals hatched in Autumn mature in the eleventh month.

Authors disagree on the length when first sexual maturity occurs. Fage (1920) and Cărașu (1952) report that it is 12 to 13 cm. Along the coast of Castellón (Western Mediterranean) the minimum length of mature fish is 111 mm for males, 117 mm for females (Andreu-Rodriguez Roda, 1951 a); on the coast of Blanes, 118 mm for males, 119 mm for females (Bas and Morales, 1954). According to Mužinić (1956) the Dalmatian anchovy matures when 9 to 10 cm long. Vidalis (1949) says that the minimum length is about 100 mm for anchovy from Greek waters and individuals 80 to 95 mm are probably mature.

3.1.3 Mating (monogamous, polygamous, promiscuous)

The anchovy is polygamous

3.1.4 Fertilization (internal, external)

External and takes place immediately upon spawning.

3.1.5 Fecundity

- Relation of gonad size and egg number to body size and to age

The relation between the sizes of the gonad and the body of the anchovy off Blanes (Western Mediterranean) given by Bas and Morales (1954) is shown in Table VI.

Slastenenko (1956) says that a female can produce as many as 40,000 eggs. Nikolski (1957) states that the average number of eggs is 13,360. Detailed investigation on the fecundity of different sizes have, however, been made by Majarova-Chugunova (1954). See Table VII.

- Coefficient of fecundity

The indices of fecundity of the Spanish anchovy from the coast of Blanes shown in Table VIII, are those given by Bas and Morales (1954).

3.1.6 Spawning

- Spawning seasons (beginning, end, peak)

The spawning season is long and not limited regularly. Its beginning, end and peak vary according to climatic conditions and areas from year to year (see Table IX).

- Number of spawnings per year, frequency

Anchovy have only one spawning season per year. But the ovarial eggs of the

Table VI

Relation of gonad size to body size from Bas and Morales (1954)

Body length	Gonad weight				Gonad length			
	Male		Female		Male		Female	
	f	M	f	M	f	M	f	M
120	12	0.34	3	0.29	12	28.5	3	27.6
130	16	0.36	20	0.59	12	28.1	18	35.8
140	20	0.53	3	0.88	15	38.2	3	37.3
150	17	0.55	1	0.67	15	34.4	-	---
160	3	0.73	-	---	3	39.3	-	---

Table VII

Number of eggs (x 1000) produced by the Black Sea anchovy (from Majarova - Chugunova, 1954)

Group	First		Second			Third			Fourth		
Number of eggs in ovarium	3 Portion		2+ Portion			2 Portion			1 Portion		
Phase of maturity	IV, V		V-VI			VI-III and VI - IV			VI-III (III) ¹		
Length of fish in cm	M	from-to	n	M	from-to	n	M	from-to	n	M	from-to
7-8	13.7	11.7-15.8	2	-	-	-	-	-	-	-	-
8-9	17.5	14.0-21.1	2	13.6	10.8-16.9	3	-	-	-	-	-
9-10	18.6	18.6	1	15.9	14.4-16.7	5	-	-	-	-	-
10-11	20.0	20.0	1	16.3	16.3	1	19.2	19.2	1	-	-
11-12	26.3	21.6-30.9	5	-	-	-	16.6	12.7-19.9	7	9.4	8.2-10.8
12-13	27.6	24.4-31.4	7	27.2	27.2	1	16.8	11.8-22.5	10	-	-
13-14	29.6	29.6	1	-	-	-	18.3	18.3	1	6.9	6.9

Table VIII

Indicia of fecundity from Bas and Morales (1954)

Length of fish in mm.	Index of fecundity	
	Frequency	Mean
120	16	2.47
130	36	3.08
140	27	3.06
160	3	2.87

Table IX
Spawning seasons

Locality	Author	S p a a w n i n g s e a s o n		
		Beginning	Peak	End
AZOV SEA	Berg et al. (1949)	June	-	August
	Slastenenko (1956)?	May	-	August
	Nikolski (1957)	end of May	-	second half of August
BLACK SEA	Soviet waters	Dekhnik (1954)	end of April or May	September October
		Vodjanitski-Kazanova (1954)	May	September
		Slastenenko (1956)	end of May	end of September
	Bulgarian waters	Gueorghiev et al. (1961)	end of May	second half of June-second half of August end of September
SEA OF MARMARA	Arim (1957)	second half of April	July	second half October
Greek waters	Vidalis (1949)	May	-	November
ADRIATIC	Split	Gamulin (1940) quoted by Vucetic (1957)	second half of April	end of August
	Lakes of Mljet Islands	Vucetic (1957)	May	September
ITALIAN WATERS	Gulf of Catania	Russo (1925) quoted by De Buen (1931)	April	August September
	Gulf of Naples	Raffaele (1888)	May	September
		Lo Bianco quoted by D'Ancona (1933)?	April	September
SPANISH MEDITERRANEAN	-	Lozano Rey (1947)	April	September
	Castellón coast	Planas and Vives (1951)	-	June-July September October
	Málaga	Miranda-Rivera (1930,1931) quoted by Andreu and Rodríguez Roda (1951)	May, June	July-August August, September
	Algerian waters	Dieuzeide et al. (1954)	April	- Second half of September
MEDITERRANEAN AREA	Page (1920)	April	June-July	September

individual fish do not mature at the same time and are not spawned all at once. By studying the size of the ovarian eggs of the Black Sea anchovy and classifying the ovaries according to a scale of maturity, Majarova-Chugunova (1954) found evidence of three different egg groups at differing degrees of maturity. They consider the first group to have been shed in June or July, the second in July or August and the third in September.

- Spawning time of day

Soviet authors are unanimous that the Black Sea anchovy spawns between midnight and 4 a.m., but, according to Vučetić (1957), it occurs between approximately 7 p.m. and 9 p.m. in the Lakes of Mljet Island (Adriatic).

- Induction of spawning, artificial fertilization

The minimum temperature for the ripening of gonads is about 13°C Fage (1920). Eggs have never been found below this temperature and if the temperature of the surface waters is high enough, anchovy tend to spawn throughout the year. This is why the spawning season is longer in southern areas than in northern ones. It can be concluded that temperature is the main factor in inducing spawning.

3.1.7 Spawning grounds

- Coastal (surface, vegetation, shore, shoal, sand, shelter); bottom
- Oceanic (surface, bottom)

Although the main concentration of eggs is in coastal waters, they are also found in oceanic waters.

The eggs float between the surface and a depth of 50 m. Dekhnik (1954) states that the main concentrations of eggs are found at 5 to 10 m depth, the number decreasing gradually down to 25 m. Below 25 m eggs are few and show signs of abnormal development.

3.1.8 Egg: structure, size, hatching type, parasites

- Structure and predators

Anchovy eggs are pelagic and differ from the eggs of most of pelagic fish in that they are ovoid rather than spherical. In the living eggs, the capsule is thin and elastic, the perivitellin space narrow, the yolk mass segmented, colorless, translucent, without oil globules.

Blastoderm forms at the lower pole of the egg and the embryo develops along the long axis and has no pigment (see Fig.2). During development, the eggs float with the long axis perpendicular to the surface of the water.

- Size

The sizes of the eggs collected from various seas are shown in Table X. By measuring the diameters of the longitudinal and transverse axes of eggs from the Turkish waters of the Black, Marmara, Aegean and Mediterranean (Gulf of Iskenderun) Seas and calculating the mean axial lengths, axial ratios, volumes of these eggs according to each sea, Demir (1959) found marked variations in egg size and shape. Fig. 3 shows that egg size or volume increases gradually and the shape becomes more spherical from the Mediterranean to the Black Sea.

Table X

Egg size of Engraulis encrasicolus

Locality		Author	Transversal axis in mm.	Longitudinal axis in mm.
BLACK SEA	Soviet waters	Berg et al. (1949)	1.2	1.9
		Vodjanitskii and Kazanova (1954)	0.8-1.2	1.5-1.9
	Turkish waters	Demir (1959)	0.66-0.95	1.05-1.58
SEA OF MARMARA		Demir (1959)	0.60-0.84	1.03-1.63
AEGEAN SEA (Turkish waters)		Demir (1959)	0.42-0.63	0.90-1.50
MEDITERRANEAN	Gulf of Naples	Raffaele (1888)	0.50-0.55	1.15-1.25
	Castellón waters	Andreu and Rodríguez Roda (1951 b)	0.51-0.61	1.22-1.50
	Gulf of Iskenderun	Demir (1959)	0.47-0.55	1.05-1.29

- Parasites and predators

All plankton-feeders prey on the eggs.

3.2 Larval history

3.2.1 Account of embryonic and juvenile life (prelarva, larva, postlarva, juvenile)

- Prelarva

The newly hatched prelarva is elongated in shape and has a long oval yolk sac. There is no mouth or pharyngeal cavity. Intestine is slender and straight. Anus is situated at the margin of embryonic fin, well to the back. Notochord is large and multicolumnar, at least partly. Pectoral fins appear as small semicircular folds on either side of the subnotochordal region and rather remote from head. There are several black chromatophores or melanophores on the ventral side of the body. The prelarval characteristics are shown in Fig. 4 which shows a somewhat advanced phase.

- Larva, postlarva

The anchovy larva and postlarva can be easily distinguished from the Clupeidae larva and postlarva because the anchovy's dorsal and anal fins are differently placed. The front of the anchovy's anal fin is ahead of the rear of the dorsal. That is, the anal would seem to overlap the dorsal, if seen from the side of the fish. In the Clupeidae, a side view would show a clear separation between the anal and dorsal fins.

In the anchovy, the tail is longer than the head; in the Clupeidae, the tail is equal in length or shorter than the head. A final difference is that anchovy larvae and postlarvae generally have fewer myotomes than Clupeidae larvae and postlarvae. Fig. 5 shows the larvae and postlarvae of the Marmara anchovy corresponding to definite development stages.

- Juvenile

The principal characteristics of the juvenile, differentiating it from the adult, are the proportion of the body and the presence of silvery lateral bands on each side of the body.

- Feeding

Prelarvae feed on yolk sac. Pavlovskaja (1955, 1961), states that prelarvae, when the mouth is formed, also consume some peridiniae and other phytoplankton organisms of a diameter of 40 to 50 μ .

The other larval stages feed on phytoplankton and zooplankton organisms. According to Karnilova (1955) the larva consumes both phytoplankton and zooplankton organisms from the time of absorption of yolk mass to a length of 5 to 6 mm. When this length is reached, zooplankton is the main food of the successive larval stages. According to Pavlovskaja (1955, 1961) when the absorption of the yolk mass is completed, larvae and juveniles separate into four length groups, each having different digestive systems and feeding habits. The larvae of the first group are 4 about 10 mm long and 70 to 80 per cent of their food is nauplii Copepoda. The length of the second group ranges from 10 to 25 mm; their main food are copepods in the states of nauplii and copepodite I. The principal food of the third group, in which the larvae are 25 to 30 mm long, is copepodite in states I-IV, adult copepods and Cladocera. In the fourth group, juveniles are 35 to 50 mm long and they feed mainly in copepods and Cladocera.

In the opinion of Nikitin (1948) quoted by Nikolski (1957) cirripedia larvae have great importance in the feeding of Black Sea anchovy juveniles.

- Rates of development and survival

- Rates of development

See Tables XI and XII and Fig. 6.

- Rates of survival

Pavlovskaja (1955) reports that the survival rates of the eggs of the Black Sea anchovy which are produced by a female are 25 to 35 percent and the survival rates of the larvae and juveniles depend specially on the quality and quantity of plankton available as food.

Karnilova (1955) recorded a survival rate between 0.1 and 9.6 percent for the number of living eggs found in one cubic meter samples of water from the Azov Sea (0 to 1 m depths) during the 1950-53 spawning seasons.

- Periods of: development and survival

From about April or May to about November or December.

- Parental care

There is no parental care.

- Parasites and predators

There is no available information on the parasites of larvae and juveniles.

The predators of larvae are all animals feeding on plankton. The predators of juveniles are large and small fish, including the anchovy itself.

Table XI

Rates of development of Engraulis encrasicolus

Author	Incubation		End of absorption of yolk mass	
	Temperature C°	Duration	Temperature C°	Duration
Raffaele (1888)	-	2 - 3 days	-	4 - 5 days
Berg et al. (1949), Cărașu (1952)	-	25-30 hours	-	-
Karlinova (1955)	22-24	38-39 hours	22-24	4 days
Vodyanitskii-Kazanova (1954)	-	24 hours	-	-
Pavlovskaja (1955)	-	-	25	30 hours
			12	120 hours
Nikolski (1957)	15	65 hours	-	-
	25-27	29 hours		
Vučetić (1957)	22	40 hours	-	-
	27	24 hours		

Table XII

Rates of development of Engraulis encrasicolus

Locality	Author	Length of larva in mm.					
		Hatching	End of absorption of yolk mass	D. rays become conspicuous	A. rays become conspicuous	Appearance of V.	Formation of scales
AZOV SEA	Kornilova (1955)	1.7	3.6-4.0	-	-	-	27.0
	Berg et al. (1949)	2.0	-	-	-	-	25.0
BLACK SEA	Vodyanitskii and Kazanova (1954)	1.58-1.60	3.0	-	-	-	-
	Pavlovskaja (1955)	1.75-1.80	4.0	6.0-7.0	6.0-7.0	-	-
SEA OF MARMARA	Demir (unpublished)	2.5-2.7	3.5-4.0	5.2-5.9	5.5-6.2	14.5-15.0	31.0-35.0
MEDITERRANEAN	Fage (1920)	3.3	4.0-5.0	-	-	15.0	35.0-40.0

3.3 Adult history

3.3.1 Longevity

Fage (1911, 1920) states that the oldest Mediterranean anchovy is two years old Majarova-Chugunova (1954), Stoianov (1961) record four-year-old anchovies in the Black Sea.

3.3.2 Hardiness

The anchovy can stand a temperature between 6 and 29°C and can accomodate itself easily to changes in salinity between 5 and 41.55 ‰ when the rise of both is gradual. Although its natural food is plankton organisms, it can also swallow fish.

3.3.3 Competitors

The main competitors are Sardina pilchardus, Sardinella aurita, Clupea sprattus, Trachurus trachurus, T. mediterraneus, Scomber scombrus.

3.3.4 Predators

Sea birds, Delphinus delphis, Tursiops tursio and all predacious fishes, especially Huso huso, Thunnus thynnus, Auxis thazard, Sarda sarda, Scomber scombrus, Belone belone, Lucioperca lucioperca, Caspialosa pontica.

3.3.5 Parasites and diseases.

In the Black Sea, nematode larvae are the well-known endoparasites of the anchovy. They infest the whole digestive system of the fish.

Reichenbach-Klinke (1958) gives the following list for the parasites of anchovy and reports that he found them (except the first) in the digestive systems and internal organs of Naples anchovies.

- TREMATODES: Aponurus lagunculus Loss
Aphanurus stossichi (Monticelli)
Metacercaria (Bucephalides?)
- CESTODES : Scolex pleuronectis Müller
(Plerocercoid of one Tetracanthidae)
- NEMATODES : Contracaecum clavatum (Rud.)
- FUNGI : Ichthyosporidium hoferi (Plehn-Mulsow)

According to Reichenbach-Klinke (1958), the last named parasite, which infest 15 percent of the fishes, causes a grave disease which kills the fish within a short time.

Although Lernaenicus encrassicola (Turton) is an ectoparasite of both anchovy and sprat (Clupea sprattus L.), there is no report that this parasite has been observed in the Mediterranean anchovy.

3.3.6 Greatest size

Mužinić (1956) has reported an Adriatic anchovy 180 mm long.

3.4 Nutrition and growth

3.4.1 Feeding (time, place, manner, season)

Since anchovy look for their food actively, feeding takes place during the daytime. According to Dimov (1961) anchovy usually feed twice during the nyctemeral cycle: in morning and evening. But a complementary feeding can occur in the middle of the day.

Feeding is much more intensive during summer than in winter. Nikolski (1957) states that the Azov Sea anchovy, after intensive summer feeding, generally interrupt feeding. When they migrate to the Black Sea in Autumn, they consume little food there until Winter's half over, from mid-Winter until Spring, the anchovy do not consume any food.

3.4.2 Food (type, volume)

According to Fage (1911), when anchovy are in surface layers, they feed on plankton organisms, preferably Copepods, Mysis and the embryos of coastal crustacean species. When they are isolated in deep waters, several diatoms and some small benthic animals in a certain quantity of clay or the fragments of annelids, Paragus and small fishes have been found among their stomach contents.

Okul (1940) quoted by Slastenenko (1955, 1956) reports this food composition in the Azov Sea: copepods 50.2 percent; phytoplankton 10.9 percent; adults and larvae of polychaets 9.5 percent; cirriped larvae 5.8 percent; fish larvae 5.1 percent and Mollusc larvae. Nikolski (1957) states that the food of the Azov Sea anchovy generally consists of copepods 60 percent; polychaets 20 percent; and cirripeds 20 percent. In the feeding of the Black Sea anchovy copepods with 35 to 55 percent, occupied the first place; decapod larvae 18 to 35 percent, the second; and mollusc larvae the third. According to Chajanova (1954), copepods, Cladocera, larvae of cirripeds, decapoda, Mysidae and Sagitta, Oikopleura, mollusc larvae constitute the main food of the Black Sea anchovy. Polychaet and fish larvae are rarely found among stomach contents. The quantity of these organisms varies with the month and locality, copepods always occupy first place in the Black Sea anchovy's diet.

Dieuzeide et al. (1954), on the other hand, say that although anchovy feed on plankton organisms, they also prey on rather large fishes.

3.4.3 Relative and absolute growth patterns and rates

See Table XIII.

3.5 Behaviur

3.5.1 Migration and local movements

The main features of the movements of anchovy are: In Spring, when the surface temperature of the water reaches a certain degree (8°C), anchovy occur in the surface layers. During this season they approach shallower waters, even penetrate into lakes, lagoons and estuaries. In Autumn, when the temperature of the water drops below 6 or 7°C, they pass into deeper and warmer waters which are 100 to 150 m deep in the Mediterranean (Fage, 1911). In the Black Sea this figure is 60 to 70 m (Berg et al., 1949) at 8°C (Nikolski, 1957).

Table XIII

Relation between age and length

Locality	Author	0	Length in mm.						
			1	1+	2	2+	3	3+	4
AZOV SEA	Berg et al. (1949)	41-77 55	71-107 87	-	97	-	-	-	-
	Nikolski (1957)	-	84	-	94	-	103	-	-
Soviet waters	Berg et al. (1949)	44-80 66	83-137 111	-	119	-	-	-	-
Eastern part (Batumi, Sukhumi)	Majarova-Chugunova (1954)	63-79	-	92-111	-	105-118	-	119-121	-
Western part (Yalta)	Majarova-Chugunova (1954)	74-84	81	109-122	114	121-128	124	128-131	129
Rumanian waters	Cărăușu (1952)	-	120-130	-	140	-	-	-	-
Bulgarian waters	Stoianov (1961)	80	99	110	118	123	128	-	138
Greek waters	Vidalis (1949)	-	90-110	130-140	-	-	-	-	-
Spanish Medite- rranean, Roussillon coast.	Page (1911, 1920)	-	120-130	-	150-170	-	-	-	-

Majarova-Chugunova (1954), who distinguish the eastern and western forms of Black Sea anchovy, give details of these movements in that sea. In the eastern Black Sea, the anchovy has its main winter quarters off the Caucasian coast. During Spring, it migrates northwards and spreads throughout the eastern oceanic water. The western form, which winters off the Crimean peninsula, migrates mainly to the northwestern area, but also spreads all over the western Black Sea during spring. At the beginning of Winter, both forms seek coastal waters. From November to January they are near the surface, but descend to 70 to 80 m later in the Winter. Between March and April, the eastern form begins the spring migration. In the northwestern Black Sea, the spring migration begins in May, with maximum intensity in June. During Summer, both forms are found above the thermocline all over the Black Sea. The main features of the Winter quarters and migration routes given by the above authors are shown in Fig. 7. The Winter quarters are characterized by special depth conditions, viz. shallow water (20 to 30 m) bordered by oceanic depths which cut into the continental terrace as gorges or submarine valleys. The current is slow in the gorges and there the anchovy aggregates. The schools vary in size and move constantly; during the day, they are at 45 to 55 m, but at night spread out and come up to 20 to 25 m.

A great number of anchovy seek Winter quarters in the Turkish waters of the Black Sea. Aasen and Akyüz (1956) recorded Winter shoals in Fatsa Bay in the eastern Black Sea. But some anchovy migrate to the Sea of Marmara during Autumn. The Spring migration toward the Black Sea generally begins at the end of April.

Soviet authors, on the other hand, say seasonal migrations take place between the Azov and Black Seas. Popova (1954) states that the Autumn migration of the Azov Sea anchovy, toward the Black Sea and its Winter quarters off the Crimean peninsula and Caucasus coast, begins in September or October. The return migration takes place in April and May. The winter quarters and migration routes of this anchovy are shown in Fig. 8.

3.5.2 Schooling

Anchovy display a schooling instinct in all phases of life.

3.5.3 Reproductive habits

See 3.1.1, 3.1.3, 3.1.4, 3.1.6, 3.1.7.

4. POPULATION (STOCK)

4.1 Structure

4.1.1 Sex ratio

According to Andreu and Rodriguez Roda (1951b) the sex ratio of Catalan anchovy by percentage is:

<u>Sex</u>	July	August	November
Males	63.8	20.8	31.3
Females	36.2	79.2	68.7

According to Crehuet-Val Cordón (1960) the sex ratio was 38.18 percent males, 61.90 percent females in 1,079 anchovies from Malaga. Mužinić (1956) records that he found 124 or 52.3 percent males, 112 or 47.3 percent females and one hermaphrodite in 227 specimens from the Adriatic. Porumb, I.I and Porumb F.I (1958) give the following sex ratios of the anchovy caught in 1954 in Rumanian littoral waters, by percentage.

	May	June	July	September	October
Males	41.86	10.97	38.46	38.09	58.33
Females	58.13	89.02	61.90	61.90	41.66

4.1.2 Age composition

Observations on age composition of anchovy stocks are relatively few. According to Page (1911), Mediterranean anchovy populations are composed of fishes one and two years old. According to Slastenenko (1956), the age composition of the Azov Sea anchovy in commercial catches, formed the following classes:

Year class	0	1	2	3
Percentage	33-88	7.58	5	0.2-0.3

Majarova-Chugunova (1954) and Stoianov (1961) give the following age composition of the Black Sea anchovy stock (by percentage) according to the averages of 1949-52 and 1952-59 respectively.

Age	1+	2+	3+	4+
Majarova-Chugunova (1954)	72.9	16.9	9.9	0.3
Stoianov (1961)	53.14 to 66.27	29.03 to 32.58	4.67 to 13.48	0.03 to 0.80

4.1.3 Size composition

According to Slastenenko (1956), the length and weight of the Azov Sea anchovy in commercial catches, vary between 80 and 110 mm, 6 and 12 g, respectively. In the Black Sea, the length composition of the anchovy catch ranges from 55 to 170 mm (Borcea) 1932; Majarova-Chugunova, 1954; Stoianov, 1961); in the Aegean Sea, from 50 to 150 mm (Vidalis, 1949; Demir, unpublished data). The length of the Israel coast anchovy is between 52 and 110 mm (Ben-Tuvia, 1953). Mužinić (1956) gives the length distribution of 237 anchovy caught along the Dalmatian coast as ranging between 80 and 180 mm with

modes 110 and 160 mm. According to Hoek (1912), quoted by Fage (1920), the length of the Naples anchovy varies from 110 to 159 mm with a mode of 120 to 129 mm. Dulzette (1938) gives the length distribution of 300 anchovy caught in the Gulf of Catania as ranging from 90 to 149 mm with a mode of 115 to 119 mm. Andreu and Rodriguez Roda (1951 a) give the following length-frequency distribution of 1867 anchovy caught in 1949-50 off the coast of Castellón:

Length (in mm)	60	70	80	90	100	110	120	130	140	150	160	170
Percentage	0.2	2.9	1.7	0.4	0.4	5.4	13.7	23.3	27.8	20.0	3.8	0.2

Crehuet and Val Cerdón (1960) give the length and weight distribution of 1,079 anchovy from Malaga. These fishes ranged from 95 to 154 mm with a mode of 120 to 124 mm and from 8.3 to 32.3 g, respectively.

4.4 Mortality, morbidity

4.4.1 Rates of mortality

According to Popova (1954), the rates of mortality of the stock of the Azov Sea anchovy vary between 40 and 80 percent, averaging 60 percent when in the Black Sea they are found spending the winter months. Mortality rates of the Black Sea anchovy off the Bulgarian coast are shown in Fig. 9.

4.6 Relation of population to community and ecosystem, biological production, etc.

Anchovy depend especially on the pelagic community. Pavlovskaja (1961) showed that in the northwestern Black Sea, a definite relation existed between the abundance of anchovy juveniles and the quantity of the zooplankton available as food.

5. EXPLOITATION

5.1 Fishing equipment

5.1.1 Fishing gear

The most important gear are: purse seines, beach seines, lampara, ring nets, trap nets, gillnets.

According to Akyüz (1959), the midwater trawl (Danish vinge-trawl) is also an important modern implement for anchovy fishing. Large quantities of anchovy can be caught by this trawl in Winter.

5.1.2 Fishing boats

No special boat types are used for anchovy

5.2 Fishing areas

5.2.1 General geographic distribution

Anchovy fisheries are located everywhere the fish is known. The important fishing areas are: the Black, Marmara, Aegean, Adriatic Seas, the South coast of Sicily (Sea of Lampedusa), the Gulf of Genova, the western coast of the Gulf of Lion, the coasts of Catalonia, Málaga and the Nemurs region of Algeria.

5.2.2 Geographical ranges (latitudes, distances from coast, etc.)

From 5 to 10 m to 15 to 20 nautical miles offshore.

5.2.3 Depth ranges

From 5 to 100 m.

5.3 Fishing seasons

5.3.3 Dates of beginning, peak and end of season

The fishing season generally lasts all the year round, with peaks either in Autumn and Spring, when migration takes place, or in Winter when the fish have moved to Winter quarters. See Table XIV.

5.3.5 Factor affecting fishing season

Water temperature seems responsible for migrations. Season peaks generally coincide with either migration times, or times when fish are in Winter quarters. Therefore, water temperature seems a valuable indicator of availability. For instance, in the Black Sea, the optimum temperature ranges of the fishing season are between 7 and 19°C. At temperatures higher than 21 or 22°C, anchovy fishing is rare.

5.4 Fishing operations and results

5.4.1 Effort and intensity

There is no sufficient data on the effort and intensity in all Mediterranean areas. But it is known that most of the effort and intensity (constituting about 30 to 40 percent of total fish catches in this area) are in the Black Sea.

5.4.3 Catches

See Table XV

Table XIV

Fishing seasons

Country	F i s h i n g s e a s o n
ALGERIA	Especially in Winter months
BULGARIA	All the year round with peaks in Spring and Autumn.
FRANCE	
Lion Gulf	All the year, with peaks in Spring and Autumn.
ITALY	Especially from March to September
RUMANIA	All the year, with peaks October-November and May.
SPAIN	
Coast of Blanes	From May to the end of August
Coast of Castellón	All the year round
Coast of Málaga	All the year round, uniformly
TURKEY	All the year, especially in Winter months
U.S.S.R.	
Black Sea	All the year, especially in Winter months
Azov Sea	From Spring to Autumn
YUGOSLAVIA	From March to October

Table XV
Annual catches in some countries

Country	Quantities (thousand metric tons)						Observations
	1955	1956	1957	1958	1959	1960	
France (incl. Algeria)	7.3	6.9	6.2	5.6	6.4	4.6	Mixed with Atlantic anchovy
Greece	1.0	5.7	6.0	9.5	7.8	-	
Italy	-	-	-	23.9	20.8	20.5	
Spain (incl. Ceuta and Melilla)	75.8	78.5	69.6	91.3	90.4	100.5	Mixed with Atlantic anchovy
Turkey	7.6	7.8	13.1	9.5	48.9	-	
U.S.S.R.	7.5	9.4	18.0	25.9	42.6	33.2	
Yugoslavia	1.5	1.9	1.6	1.2	1.4	1.0	

SYNOPSIS OF BIOLOGICAL DATA ON ANCHOVY - N. DEMIR

6. REFERENCES

- Aasen, O. and E. Akyüz, Fishery investigations in Turkish Black Sea waters with
1956 special reference to anchovy. Rep. Fish.Res.Cent. Istanbul (Mar.Res.)
1 (7): 39 p.
- Akşiray, F., Türkiye deniz balıkları tayin anahtarı. Istanbul, Hydrobiological
1954 Research Institute, 177 p.
- Akyüz, E.F., On the use of midwater trawls for anchovy in the Black Sea. Modern
1959 fishing gear of the world, ed. by H. Kristjonsson, London, Fishing
News (Books) Ltd., pp. 357-8.
- *Alexandrov, A., Anchois de la mer d'Azoff et de la Mer Noire, leur origine et
1927 indications taxonomiques. Rep. sci. Sta. Fish. Kertch, (1)
- Andreu, B. and J. Rodriguez Roda, La pesca marítima en Castellón.- Rendimiento por
1951a unidad de esfuerzo (1945-1949) y consideraciones biométricas de las espe-
cies de interés comercial. Pub.Inst.Biol.apl. Barcelona, (8): 233-77.
- Andreu, B. and J. Rodriguez Roda, Estudio comparativo del ciclo sexual, engrasamien-
1951b to y repleción estomacal de la sardina, alacha y anchoa del mar Catalán,
acompañado de relación de pescas de huevos planctónicos de estas especies.
Publ.Inst.Biol.apl.Barcelona, (9): 193-232.
- *Arim, N., Marmara ve Karadenizdeki bazı kemikli balıkların (teleost'ların) yumurta
1957 ve larvalarının morfolojileri ile ekolojileri. Hidrobiol., Istanbul, A4
(1-2).
- Aurich, H.J., Engraulis encrasicolus (L.)
- Bas, C. and E. Morales, Algunos datos para el estudio de la biología de la anchoa
1954 Engraulis encrasicolus L. de la Costa Brava. Publ.Inst.Biol.apli.Barce-
lona, (16): 53-70.
- Ben-Tuvia, A., Mediterranean fishes of Israel. Bull.Sea Fish.Res.Lta., Haifa, (8):
1953 40 p.
- Berg, L.S., Classification of fishes both recent and fossil. Frud.zool.Inst.Akad.
1947 Nank SSSR, 5(2)
- Berg, L.S. et al (Ed), Promyslovye ryby SSSR. Morkva, Pischepromizdat, 787 p.
1949
- *Borcea, I., Observations sur les poissons migrants dans les eaux Roumaines de la
1929 Mer Noire. Ann.sci.Univ.Jassy, 15.
- *Borcea, I., Nouvelles observations sur les migrations et sur le période de la ponte
1932 des espèces de poissons migrants de la Mer Noire. Ann.sci.Univ.Jassy, 17
- Cărbăușu, S.I., Tratat de ichtiologie, Bucharest, Edit. Academiei Republicii Populare
1952 Române, 802 p.
- Carus, J.V., Prodromus faunae mediterraneae, 2. Stuttgart, 854 p.
1893

- *Chaianova, L.A. Pitanie Chernomorsko i kamsy. Frud.vsesoinz.nauch.-
1954
- *Crehuet, R.F. and M.J. Val Cordon, Estudio preliminar sobre la biométrica, biología
1960 y variación del contenido graso del boquerón (Engraulis encrasicolus)(L.))
de Málaga. Bol.Inst.esp.Oceanogr., (99)
- D'Ancona, U., Uova, larvae e stadi giovanili di teleostei, Clupeidae. Monogr.Stazool.
1931 Napoli, (38): 176 p.
- De Buen, F., Catálogo ictiológico del Mediterráneo español y de Marruecos. Madrid,
1926 221 p.
- *De Buen, F., Clupéidés et leur pêche, deuxième note. Rapp.Comm.int.Mer Médit., 6
1931
- *De Buen, F., Clupéidés et leur pêche, troisième note. Rapp.Comm.int.Mer Médit., 7
1932
- *Dekhnik, T.V., Razmnazhenie khamsy i kefali v chernom-lore. Frud.vsesoinz.nauch.-
1954 issled.Inst.morsk.ryb.Khoz., 28
- *Demir, N., Notes on the variations of the eggs of anchovy (Engraulis encrasicolus
1959 Cuv.) from Black, Marmara, Aegean and Mediterranean seas. Hidrobiol.
Istanbul, B4 (4)
- Devedjian, K., Pêche et pêcheries en Turquie. Istanbul, 480 p.
1926
- Dieuzeide, R., M. Novella and J. Roland., Catalogue des poissons des côtes algé-
1954 riennes. Bull.Sta.Acquic.Pêche, Castiglione, (5): 258 p.
- Dimov, I., Sur le rythme nyctéméral de la nutrition chez l'anchois (Engraulis
1961 encrasicolus ponticus Alex.) et le sprat (Sprattus sprattus sulinus Ant.)
Izv.nauch.- issled.Inst.Rib.Varna, (1): 79-94.
- Dulzetto, F., Sui caratteri biometrici dell'acciuga "Engraulis encrasicolus L."
1938 del Golfo di Catania. Boll.Pesca Piscic.Idrobiol., 14 (2): 180-209.
- Einarsson, H. and N. Görtürk. Abundance and distribution of eggs and larvae of the
1960 anchovy (Engraulis encrasicolus ponticus) in the Black Sea. Hidrobiol.,
Istanbul, B5: (1-2): 72-94.
- Faber, G.L., The fisheries of the Adriatic and the fish thereof, London, B. Quaritch,
1883 292 p.
- Fage, L., Recherches sur la biologie de l'anchois (Engraulis encrasicolus Linné),
1911 races-age-migrations. Ann.Inst.océanogr., Paris, 2(4): 140 p.
- Fage, L., Engraulidae, Clupeidae. Rep.Danish oceanogr.Exped.Medit., 2(A9): 136 p.
1920
- * Fage, L., Engraulidae. Faune ichthyol.Atlant.N., (4)
1930
- Fage, L., La ponte et les races locales de l'anchois de la Méditerranée (Engraulis
1937 encrasicolus L.). Rapp.Comm.int.Mer Médit., 10: 67-71.

- GFCM, Catalogue of the names of fishes of commercial importance in the Mediterranean.
1960 Rome. FAO, 250 loose sheets.
- FAO, Yearb.Fish.Statist. F.A.O. (Prod.), 12(1960)
1960
- Fowler, H.W., The marine fishes of West Africa, Based on the collection of the
1936 American Museum Congo expedition, 1909-1915. Bull.Amer.Mus.Nat.Hist., 70(1):
605 p.
- *Gueorghiev, J.M. K. Alexandrova and D.C. Nikolov. Distribution quantitative des
1961 oeufs et des larves de l'anchois (Engraulis encrasicolus ponticus Alex.)
du saurel (Trachurus mediterraneus ponticus Aleev) et du rouget (Mullus
barbatus ponticus Essipov) devant le littoral bulgare de la Mer Noire.
Izv.nauch.- izsled.Inst.Rib.Varna, (1)
- *Günther, A., Catalogue of the fishes in the British Museum, vol.7, London.
1868
- *Holt, E.W.L., Recherches sur la reproduction des poissons osseux principalement dans
1899 le Golfe de Marseille, Memoire 2. Ann.Mus.Hist.Nat.Marseille, Zool., 5
- Karnilova, V.P. Trud.azov.-chernomosrk.nauch.-issled.Inst.morsk.ryb.Khoz., (16)
1955 Nabludenie za rostom lichinok i malodi Azovskoi khamsy v 1953 g.
- Kyle, H.M. and E. Ehrenbaum. Vertebrata, Die Fische der Nord-und Ostsee. 12 (c-h).
1927 Tierwelt N.- u.Ostsee.
- *Larraneta, M.G., Sur la formule vertébrale de quelques poissons commerciaux des cô-
1958 tes de Castellón. Rapp.Comm.int.Mer Médit., 14.
- Lozano Rey, L., Peces ganoideos y fisóstomas. Mem.R.Acad.Madri., 9:840 p.
1947
- Lübbert, H. and E. Ehrenbaum, Handb.Seefisch.Nordeurop., 2
1936
- Maierova, A.A. and N.I. Chugonova, Biologiya, raspredelenie i otsenka zapasa
1954 chernomorskoj khamsy. Trud.vsesoinz.nauch.- issled.Inst.morsk.ryb.Khoz.,
28:5-33.
- Moreau, E., Histoire naturelle des poissons de la France. Paris, 3:697 p.
1881
- Mužinić, R., Quelques observations sur la sardine, l'anchois et le maquereau des
1956 captures au chalut dans l'Adriatique. Acta adriat., 7(13): 39 p.
- Nikolskii, G.V., (Hentschel, F.Y. Fransl.) Spezielle Fischkunde, Berlin, Deutscher
1957 Verlag der Wissenschaften, 632 p.
- Ninni, E., Catalogo dei pesci del mare adriatico, Venezia. 271 p.
1912
- Palombi, A. and M. Santarelli, Gli animali commestibili dei mari d'Italia. Milano.
1953 U. Hoepli, 349 p.

- *Pavlovskaja, R.M., Vyzhivanie chernomorskoj khamsy na rannikh etapakh razvitiia.
1955 Frud.azov.-cherno-morsk nauch.- issled.Inst.morsk.ryb.Khoz.
- *Pavlovskaja, R.M., La survie des larves de l'anchois de la Mer Noire en fonction de
1961 leur alimentation. Rapp. Comm.int.Mer Médit., 16.
- Planas, A. and F. Vives, sobre la puesta de la anchoa (Engraulis encrasicholus (L.)
1951 en el levante español. Publ.Inst.Biol.apli.Barcelona, (9):119-30)
- *Popiel, Y., Clupeidae, Engraulidae, Scombridae, Klucze do Oznaczania kregowcow Polski.
1962 Polok.Akad.Nauk.Zakl.Zool.Syst.Krakow, (1)
- Popova, V.P., Chernomorskii period zhizni azovskoi khamsy. Frud.vsesoizn.nauch.-
1954 issled.Inst.morsk.ryb.Khoz., (28)
- Porumb, I.I. and F.I. Porumb, Contributions à l'étude de la biologie des poissons
1958 pélagiques en face du littoral roumaine de la Mer Noire. Anl.stünt.Univ.
Al.Y.Cuza, 4:77-101.
- *Raffaele, F., Le uova galleggianti e le larve dei teleostei del Golfo di Napoli.
1888 Mitt.zool.Sta Neapel, 8(1).
- *Reichenbach - Klinke, H.H., les parasites de la sardine (Sardina pilchardus Walb.)
1958 et de l'anchois (Engraulis encrasicholus Rond.). Rapp.Comm.int.Mer Médit.,
14.
- Scott, T. and A. Scott, The British parasitic Copepoda. London vol.1:256 p.,
1913 vol 2: plates.
- *Slastenenko, E.P., Les poissons de la Mer Noire et de la Mer d'Azov. Tra.Sta.zool.
1941 marit. Regele Ferdinand I Agigéa, (3).
- Slastenenko, E.P. Karadeniz havzasi baliklari (the fishes of the Black Sea basin),
1956 Istanbul, 711 p.
- *Stoianov, S., Etat du stock de l'anchois de la Mer Noire, pêché devant le littoral
1961 bulgare pendant la période 1952-1960. Izv.nauch.-izsled.Inst.Rib.Varna,
(1).
- Thompson D'Arcy W., A glossary of Greek Fishes, London, Oxford University Press,
1947 302 p.
- Vidalis, E., Contribution in the biology of the anchovy in Greek waters. Prakt.Hellen.
1949 hydrobiol.Inst., 3(2): 41-70.
- *Vodianitskii, V.A. and I.I. Kazanova, Opredeletel pelagicheskii'lichinok ryb chernogo
1954 Moria. Frud.vsesoizn.nauch.- issled.Inst.morsk.ryb.Khoz., 28
- Vučetić, T., Quelques observations sur l'écologie de la ponte de l'anchois (Engraulis
1957 encrasicholus (L.)) dans les lacs de l'Ile de Mljet. Proc.gen.Fish.Coun.Medit.,
4: 227-33.
- *Whitehouse, R.H., Report on fish eggs and larvae taken during 1931. Notes Fish.Res.
1933 Dir., Cairo, (4).

Appendix

FIGURES




Fig. 1 - Sprat, from the Gulf of Erdek, Sea of Marmara




Fig. 2 - Eggs of Engraulis encrasicolus from the Sea of Marmara (preserved material), A=0.60 x 1.35 mm, B=0.65 x 1.25 mm.










Fig. 3 - Comparison of the eggs in mean size and shape originating from the Mediterranean (1), Aegean (2), Marmara (3) and Black (4) Seas. a= Longitudinal axis, b= transversal axis, the measurement are shown by unities (38 unities = 1 mm). From Demir (1959).



Fig.4 - Prelarva (2.9 mm) of Engraulis encrasicolus from the Sea of Marmara (preserved material)



H




Fig. 5 - Larvae and postlarvae (A = 3.3 mm., B = 6 mm., C = 9 mm., D = 12 mm., E = 18 mm., F = 24 mm., G = 27 mm., H = 31.5 mm metamorphosed specimen) of Engraulis encrasicolus from the Sea of Marmara (preserved material).

